

General Solution for Self-Gravitating Spherical Null Dust

Aleksandar Miković ^{1 2 3}

Departamento de Física Teórica and IFIC, Centro Mixto Universidad de Valencia-CSIC, Facultad de Física, Burjassot-46100, Valencia, Spain

ABSTRACT

We find the general solution of equations of motion for self-gravitating spherical null dust as a perturbative series in powers of the outgoing matter energy-momentum tensor, with the lowest order term being the Vaidya solution for the ingoing matter. This is done by representing the null-dust model as a 2d dilaton gravity theory, and by using a symmetry of a pure 2d dilaton gravity to fix the gauge. Quantization of this solution would provide an effective metric which includes the back-reaction for a more realistic black hole evaporation model than the evaporation models studied previously.

PACS numbers: 04.20.Jb, 04.40.-b, 04.70.Dy

¹On leave from Institute of Physics, P.O. Box 57, 11001 Belgrade, Yugoslavia

²Supported by Comisión Interministerial de Ciencia y Tecnología

³E-mail: mikovic@lie1.ific.uv.es

Two-dimensional (2d) dilaton gravity theories have turned out to be very useful toy models of black hole formation and evaporation [1]. They are also relevant for four-dimensional (4d) black holes, since a spherically symmetric scalar field collapse can be described by a 2d dilaton gravity model

$$S_0 = \frac{1}{2} \int d^2x \sqrt{-g} \left[e^{-2\Phi} \left(R + 2(\nabla_\mu \Phi)^2 + 2e^{2\phi} \right) - \frac{G}{2} e^{-2\Phi} (\nabla_\mu f)^2 \right] \quad , \quad (1)$$

where G is the Newton constant and the 4d line element ds_4 is related to the 2d line element ds by

$$ds_4^2 = ds^2 + e^{-2\Phi} d\Omega^2 \quad . \quad (2)$$

R is a 2d scalar curvature associated with a 2d metric $g_{\mu\nu}$, ∇_μ is the corresponding covariant derivative, Φ is the dilaton field, f is the scalar field and $d\Omega$ is a two-sphere line element. Quantization of (1) would provide us with a semi-classical metric which would include the back-reaction. However, the progress is hindered by the absence of explicit solutions of the classical equations of motion. In the case of 2d black holes described by the CGHS model [2], the analog of (1) is exactly solvable, and by quantizing the solution, one can obtain an effective semi-classical metric up to any finite order in matter loops [3, 4, 5]. Since the matter in the CGHS model is 2d conformally coupled, this motivates us to consider a modification of (1)

$$S = \frac{1}{2} \int d^2x \sqrt{-g} \left[e^{-2\Phi} \left(R + 2(\nabla_\mu \Phi)^2 + 2e^{2\phi} \right) - \frac{G}{2} (\nabla_\mu f)^2 \right] \quad , \quad (3)$$

so that f will obey a free-field equation of motion in the conformal gauge, and consequently the general solution could be found more easily. Action (3) describes the dynamics of spherically symmetric self-gravitating null-dust (SSND). For purely ingoing matter, the solution of the equations of motion is given by the Vaidya metric [6]

$$ds^2 = - \left(1 - \frac{2m(v)}{r} \right) dv^2 + 2dvdr \quad , \quad (4)$$

where $r = \exp(-\Phi)$, $G = 1$ and

$$\frac{dm(v)}{dv} = T_{vv}(v) = \frac{1}{2} (df/dv)^2 \quad . \quad (5)$$

An analytic model of black hole evaporation based on the quantization of the Vaidya solution has been studied in [7], and a qualitative agreement with the numerical results of [8] has been found. However, in order to fully take into account the back-reaction in the analytic approach, one needs the most general solution for which the outgoing matter is also present [9, 7].

In order to find the general solution of the SSND model, we rewrite the action (3) as

$$S = \frac{1}{2} \int d^2x \sqrt{-\tilde{g}} \left[\tilde{R}\phi + V(\phi) - \frac{1}{2}(\tilde{\nabla}_\mu f)^2 \right] \quad , \quad (6)$$

where $\phi = e^{-2\Phi} = r^2$, $\tilde{g}_{\mu\nu} = r g_{\mu\nu}$ and $V = 2/r$. We do this in order to establish the connection with a generic 2d dilaton gravity model, which can be represented by the action of the form (6). For example, the CGHS model is given by $V = 4\lambda^2$, where λ is a 2d cosmological constant. The equations of motion are given by

$$\nabla_\mu \nabla_\nu \phi - \frac{1}{2} g_{\mu\nu} V = -T_{\mu\nu} \quad , \quad (7)$$

$$R + \frac{dV}{d\phi} = 0 \quad , \quad \square f = 0 \quad , \quad (8)$$

where we have omitted the tildas, $\square = g^{\mu\nu} \nabla_\mu \nabla_\nu$ and

$$T_{\mu\nu} = \partial_\mu f \partial_\nu f - \frac{1}{2} g_{\mu\nu} (\nabla_\mu f)^2 \quad .$$

In the conformal gauge $d\tilde{s}^2 = -e^{2\rho} dx^+ dx^-$ one obtains

$$\partial_+ \partial_- \phi = -\frac{1}{4} V e^{2\rho} \quad , \quad (9)$$

$$\partial_+^2 \phi - 2\partial_+ \rho \partial_+ \phi = -(\partial_+ f)^2 \quad , \quad (10)$$

$$\partial_-^2 \phi - 2\partial_- \rho \partial_- \phi = -(\partial_- f)^2 \quad , \quad (11)$$

$$\partial_+ \partial_- \rho = -\frac{1}{8} \frac{dV}{d\phi} e^{2\rho} \quad , \quad (12)$$

and $\partial_+ \partial_- f = 0$. We want to solve the system (9-12) in analogy to the CGHS case, where a gauge $\rho = 0$ can be chosen. This is possible since $\partial_+ \partial_- \rho = 0$ is a consequence of the equations of motion. In the SSND case ρ is not a free field, so that we have to find an appropriate modification. We will use as the starting point the free-field currents for the $f = 0$ case [11]

$$j_1 = \int \frac{d\phi}{2E + J(\phi)} \quad , \quad (13)$$

$$\tilde{j}_2 = \log(\nabla\phi)^2 \quad , \quad (14)$$

where $2E = (\nabla\phi)^2 - J(\phi)$, $dJ/d\phi = V$ and E is a constant of motion. j 's satisfy

$$\square j_1 = 0 \quad , \quad \square \tilde{j}_2 + R = 0 \quad . \quad (15)$$

By going into the conformal gauge one can see that the equations (15) imply that j_1 and $j_2 = \tilde{j}_2 - 2\rho$ are free fields. In the SSND case, when $f = 0$ one has $2j_1 =$

$r + 2M \log |r/2M - 1|$ and $j_2 = \log |r - 2M| - 2\rho$, where $M = -E/4$ is the black hole mass. The Schwarzschild solution is obtained for

$$2j_1 = \frac{1}{2}(v - u) \quad , \quad (16)$$

$$j_2 = 0 \quad . \quad (17)$$

The gauge choice (16) gives the familiar relation between r and u, v coordinates, while the gauge choice (17) is equivalent to

$$ds^2 = -\frac{e^{2\rho}}{r} du dv = -(1 - 2M/r) du dv. \quad (18)$$

Note that the gauge choice $j_2 = 0$ gives a relation between ρ and ϕ , which is of the type we are looking for, and hence we will concentrate on it.

When the matter is added, the relations (15) change as

$$\begin{aligned} \square j_1 &= \frac{\partial^2 j_1}{\partial E^2} T^{\mu\sigma} T_\mu{}^\nu \nabla_\sigma \phi \nabla_\nu \phi + \left(\frac{\partial j_1}{\partial E} - \frac{4}{(\nabla \phi)^4} \right) T_{\mu\nu} \nabla^\mu \phi \nabla^\nu \phi \quad , \\ \square \tilde{j}_2 + R &= -\frac{4}{(\nabla \phi)^4} [T^{\mu\sigma} T_\mu{}^\nu \nabla_\sigma \phi \nabla_\nu \phi + V(\phi) T_{\mu\nu} \nabla^\mu \phi \nabla^\nu \phi] \\ &\quad + \frac{2}{(\nabla \phi)^2} T_{\mu\nu} \nabla^\mu \phi \nabla^\nu \phi \quad , \end{aligned} \quad (19)$$

so that j 's are not free fields any more. The form of (19) and our remark about j_2 suggest that we look for a free field of the form

$$j = j_2 + X \quad , \quad (20)$$

where X is to be determined from (19). This gives

$$\partial_+ \partial_- X = - \left[\frac{T_{++}}{(\partial_+ \phi)^2} + \frac{T_{--}}{(\partial_- \phi)^2} \right] \partial_+ \partial_- \phi \quad . \quad (21)$$

Equation (21) is valid for any potential V , and can be solved as

$$X = X_0 + \log \left| \frac{\partial_- \phi}{\partial_+ \phi} \right| + 2 \int dx^- \frac{T_{--}}{\partial_- \phi} \quad , \quad (22)$$

where X_0 is a free-field solution.

In the SSND case $\phi = r^2$, $x^+ = v$, $x^- = u$, and we choose $X_0 = 0$, which fixes the gauge. It is convenient to use a generalized form of the Vaidya metric [10]

$$ds^2 = -e^{2\psi} F dv^2 + 2e^\psi dr dv \quad , \quad (23)$$

which can be related to the conformal form of the metric via $ds^2 = -C(u, v)dudv$ and $C = e^{2\rho}/r$. Then the gauge choice $j = 0$ is equivalent to

$$\psi = 2 \int du \frac{T_{uu}}{\partial_u r^2} \quad , \quad (24)$$

where $T_{uu} = \frac{1}{2}T_{--}$. Note that we have not specified the limits of u integration in (24), which means that a constant of integration will occur. Only when this constant is specified, the gauge will be completely fixed.

The equations which determine r are (9) and (10), while (11) and (12) are the consistency conditions for the gauge choice, which are satisfied by construction. Equation (9) becomes

$$\partial_u \partial_v r^2 = e^\psi \partial_u r \quad , \quad (25)$$

while (10) gives

$$\partial_v^2 r^2 - 2\partial_v \rho \partial_v r^2 = -(\partial_v f)^2 \quad , \quad (26)$$

where ρ is determined from the gauge choice as

$$2\rho = \log | -\partial_u r^2 | + \psi \quad . \quad (27)$$

Equation (11) follows from (27) and (24), while the equation (12) follows from (27), (25) and (24). By integrating (25) with respect to u , we obtain

$$\partial_v r = \frac{1}{2} \left(1 - \frac{2m(v)}{r} \right) + \frac{1}{2r} \int du (e^\psi - 1) \partial_u r \quad . \quad (28)$$

$m(v)$ is a constant of integration, and can be determined from (26). By inserting (28) into (26), and by using (27), we get

$$T_{vv} = \frac{dm}{dv} - \frac{1}{2}(1 - e^\psi - 2r\partial_v \psi)\partial_v r - \frac{1}{2}\partial_v \int du (e^\psi - 1)\partial_u r \quad . \quad (29)$$

When $\psi = 0$, (28) yields $F = 1 - 2m/r$ and (29) gives the relation (5), so that one recovers the Vaidya solution. When $\psi \neq 0$, the equations look difficult. However, their form is such that a perturbative solution in T_{uu} can be easily found. Let us introduce an expansion parameter ϵ by replacing T_{uu} with ϵT_{uu} . Then we will seek a solution in the form

$$r = r_0 + \epsilon r_1 + \epsilon^2 r_2 + \cdots \quad . \quad (30)$$

The expansion (30) then implies

$$\psi = \epsilon \psi_1 + \epsilon^2 \psi_2 + \cdots \quad , \quad (31)$$

where

$$\psi_1 = \int du \frac{T_{uu}}{r_0 \partial_u r_0} \quad , \quad \psi_2 = - \int du T_{uu} \frac{\partial_u(r_1 r_0)}{(r_0 \partial_u r_0)^2} \quad , \quad \dots \quad . \quad (32)$$

In general one should also take

$$m(v) = m_0(v) + \epsilon m_1(v) + \epsilon^2 m_2(v) + \dots \quad , \quad (33)$$

although it is possible that the series in (33) gets truncated, like in the example we consider in this paper, where $m = \epsilon m_1$. By inserting the expansions (30), (31) and (33) into the equation for r (28), we get an infinite hierarchy of equations

$$\begin{aligned} \partial_v r_0 &= \frac{1}{2} \left(1 - \frac{2m_0}{r_0} \right) \quad , \\ \partial_v r_1 &= \frac{m_0}{r_0^2} r_1 - \frac{m_1}{r_0} + \frac{1}{2r_0} \int du \psi_1 \partial_u r_0 \quad , \\ \partial_v r_2 &= \frac{m_0}{r_0^2} r_2 - \frac{m_2}{r_0} - \frac{r_1}{r_0} \partial_v r_1 + \frac{1}{2r_0} \int du \left[\left(\frac{1}{2} \psi_1^2 + \psi_2 \right) \partial_u r_0 + \psi_1 \partial_u r_1 \right] \quad , \quad \dots \quad . \end{aligned} \quad (34)$$

The system (34) can be solved, since for every n the equation for r_n does not involve r_k with $k > n$ and each equation is a first order linear differential equation for r_n , except for $n = 0$, which is a non-linear first order differential equation. Therefore by starting from $n = 1$, one can write an explicit solution for any r_n in terms of r_0 and T_{uu} . At each step several integration constant $C(u)$ and $C(v)$ will arise (due to u and v integrations), and these constants can be determined from the constraint equation (29), exactly as in the $T_{uu} = 0$ case. This can be done because the equation (29) also decomposes into an infinite hierarchy of equations under the expansions (30), (31) and (33)

$$\begin{aligned} T_{vv} &= dm_0/dv \quad , \\ 0 &= dm_1/dv + \frac{1}{2}(\psi_1 + 2r_0 \partial_v \psi_1) \partial_v r_0 - \frac{1}{2} \partial_v \int du \psi_1 \partial_u r_0 \quad , \\ 0 &= dm_2/dv + \frac{1}{2}(\psi_2 + \frac{1}{2} \psi_1^2 + 2r_0 \partial_v \psi_2 + 2r_1 \partial_v \psi_1) \partial_v r_0 + \frac{1}{2}(\psi_1 + 2r_0 \partial_v \psi_1) \partial_v r_1 \\ &\quad - \frac{1}{2} \partial_v \int du [(\psi_2 + \frac{1}{2} \psi_1^2) \partial_u r_0 + \psi_1 \partial_u r_1] \quad , \quad \dots \quad . \end{aligned} \quad (35)$$

The equations (35) will also determine the $m_n(v)$, provided we fix the integration constants $\psi_n(v)$ which appear in (32). These are related to the complete specification of the gauge, or equivalently, to the choice of the v coordinate, since a coordinate change $v = v(\tilde{v})$ in (23) gives

$$\tilde{\psi} = \psi + \log \frac{dv}{d\tilde{v}} \quad . \quad (36)$$

At the end, one sets $\epsilon = 1$ and writes the solution as

$$r = r_0 + r_1 + r_2 + \cdots \quad , \quad \psi = \psi_1 + \psi_2 + \cdots \quad . \quad (37)$$

These general features can be nicely illustrated on the example of a white hole emitting a shock-wave. This is simply a shock-wave Vaidya solution where the u and v coordinates are interchanged. The corresponding spacetime is described by the Penrose diagram of Fig.1. In this case $T_{uu} \neq 0$, and one can find an exact solution




Figure 1: Penrose diagram of a white hole emitting a shock-wave.

for $r = r(u, v)$, so that the expansions (30) and (31) and the corresponding equations (34) and (35) can be checked. By taking $T_{uu} = M\delta(u - u_0)$, we get

$$ds^2 = \begin{cases} -du^2 - 2dudr & u > u_0 \\ -(1 - 2M/r)du^2 - 2dudr & u < u_0 \end{cases} . \quad (38)$$

This can be rewritten as

$$ds^2 = \begin{cases} -dv^2 + 2dvdr & u > u_0 \\ -Fe^{2\psi}dv^2 + 2e^\psi dvdr & u < u_0 \end{cases} , \quad (39)$$

where $F = 1 - 2M/r$,

$$\psi = -\log \left| 1 - \frac{4M}{v - u_0} \right| \quad (40)$$

and $r = \frac{1}{2}(v - u)$ for $u > u_0$, while for $u < u_0$

$$r + r_s \log |r/r_s - 1| = \frac{1}{2}(v - u) + r_s \log |(v - u_0)/2r_s - 1| \quad , \quad (41)$$

where $r_s = 2M$. The form of the solution, given by (40) and (41), is such that

$$r = \sum_{n \geq 0} r_s^n \tilde{r}_n \quad , \quad \psi = \sum_{n \geq 1} r_s^n \tilde{\psi}_n \quad , \quad (42)$$

which is of the form (37). More exactly, one can show that $r_n = r_s^n \tilde{r}_n$ and $\psi_n = r_s^n \tilde{\psi}_n$ satisfy the equations (34) and (35) with $m = m_1 = M$. Also, starting from $n = 2$, non-trivial integration constants $\psi_n(v)$ appear.

In conclusion we can say that we have found a useful form of the general solution, given by the expansions (37) and the equations (34) and (35). The expansions (37) are clearly in powers of the outgoing energy-momentum tensor, and by truncating them at finite n we obtain an explicit perturbative solution. This form of the solution can be used to construct an effective quantum metric in the approximation of a finite number of matter loops via the method of quantization of the classical solution [3, 4, 5, 7]. The corresponding construction is going to be more involved than in the 2d case [3, 4, 5] or the Vaidya case [7], since the relation between the metric and the dilaton (radius) is more complicated. For example, for the one-loop approximation one would take $r = r_0 + r_1$ and $\psi = \psi_1$, with $T_{\mu\nu}$ replaced by $\langle T_{\mu\nu} \rangle$ evaluated in an appropriate quantum state. Note that in the one-loop case one does not have to truncate the expansions (37) at $n = 1$. By including the higher-order terms, one extends the validity of the one-loop approximation to a smaller radius.

Our solution can also serve as a good starting point for finding approximate analytic solutions for the more realistic collapse described by the action (1).

Acknowledgements

I would like to thank Jose Navarro-Salas and Javier Cruz for useful discussions.

References

- [1] A. Strominger, *Les Houches Lectures on Black Holes*, Santa Barbara preprint (1995), hep-th/9501071
- [2] C. G. Callan, S. B. Giddings, J. A. Harvey and A. Strominger, Phys. Rev. D 45 (1992) R 1005.
- [3] A. Miković, Phys. Lett. B 355 (1995) 85
- [4] A. Miković, Class. Quant. Grav. 13 (1996) 209
- [5] A. Miković and V. Radovanović, Nucl. Phys. B 481 (1996) 719
- [6] P.C. Vaidya, Proc. Indian Acad. Sci. A33 (1951) 264

- [7] J. Cruz, A. Miković and J. Navarro-Salas, Phys. Lett. B 395 (1997) 184
- [8] R. Parentani and T. Piran, Phys. Rev. Lett. 73 (1994) 2805.
- [9] W.A. Hiscock, Phys. Rev. D 23 (1981) 2813
- [10] J.M. Bardeen, Phys. Rev. Lett. 46 (1981) 382
- [11] J. Cruz, J. Navarro-Salas, M. Navarro and C.F. Talavera, "*Symmetries and Black Holes in 2D Dilaton Gravity*", Univ. of Valencia preprint (1996), hep-th/9606097; "*Conformal and non-conformal symmetries in 2D dilaton gravity*", to appear in Phys. Lett. B